



Australian
Rheumatology
Association

PATIENT INFORMATION ON ALLOPURINOL

(Brand names: Allohexal, Allosig, Progot, Zyloprim)

This information sheet has been produced by the Australian Rheumatology Association to help you understand the medicine that has been prescribed for you. It includes important information about:

- How you should take your medicine;
- What are the possible side effects;
- What tests you may have to monitor your condition and to detect unwanted effects; and
- Other precautions you should take.

Please read it carefully and discuss with your doctor.

What is allopurinol?

Allopurinol (brand names Allohexal, Allosig, Progot, Zyloprim) is a medicine used to treat gout, which is a type of arthritis caused by a build up of uric acid crystals in the joints.

Uric acid is produced normally by cells, but in gout the body does not flush it out fast enough. Allopurinol works by reducing the amount of uric acid made by cells.

Allopurinol does not treat the pain or inflammation of an 'attack' of gout, but it is taken on a long term basis to prevent further attacks. The treatment also helps to reduce the risk of your joints being permanently damaged.

In addition to treating gout, allopurinol is used to treat certain types of kidney stones. It is also used in patients receiving cancer chemotherapy to prevent high uric acid levels caused by the release of uric acid from the dying cancer cells.

For more information about GOUT see the Arthritis Australia website
www.arthritisaustralia.com.au/index.php/arthritis-information/information-sheets.html.

IMPORTANT INFORMATION

Once you are prescribed allopurinol, it is very important that you **do not stop taking it** unless advised by your doctor or unless side effects develop. If allopurinol treatment is stopped there is a high risk that your gout may worsen.

You should continue to take it even if you do not have symptoms.

You should also continue to take it during an attack of gout.

What benefit can you expect from your treatment?

Following a diagnosis of gout, allopurinol may be given on a long term basis to **prevent** further attacks. It is not used to treat a sudden attack and it is not normally started during a sudden attack.

There are two important things to keep in mind when you start taking allopurinol:

- it may take several weeks to reduce the level of uric acid, so you may continue to have gout attacks for some time; and
- sometimes, starting allopurinol or increasing the dose can actually cause an attack of gout.

This does not mean the medication is not working, so keep taking it during such attacks together with any other medicine your doctor may recommend to manage the pain. Medicines such as colchicine or anti-inflammatory drugs may be recommended before or at the same time allopurinol is started to reduce the risk of a gout attack. Your doctor will advise you about how these medicines should be taken.

What happens if you have a gout attack?

Allopurinol is not a pain reliever. You should continue to take allopurinol during an attack, but your doctor will also recommend medicines to treat the pain and inflammation. These may include paracetamol, colchicine, anti-inflammatory drugs such as naproxen (Naprosyn), ibuprofen (Brufen/Nurofen), indomethacin (Indocid) or steroids such as prednisolone.

It is a good idea to plan with your doctor what to do if a gout attack occurs and to have symptom controlling medicine ready to use if you need to. It is also important that you tell your doctor if your condition persists or worsens.

How is allopurinol taken?

Allopurinol is taken by mouth as a tablet. It is usually taken once a day. If your dose is more than 300 mg day, you may need to take it a number of times a day.

It should be taken after food to reduce stomach upset. It should also be taken with plenty of water.

For greatest benefit, allopurinol should be taken regularly. To help you remember, take it at the same time(s) each day. If you forget to take a dose, there is no need to double the dose at the next scheduled time.

What is the dosage?

Tablets come in 100mg or 300mg strengths. Treatment may start with a small dose, increasing over time to between 100mg and 300mg per day. Occasionally higher doses are used. If you have kidney problems the dose is likely to be lower.

Are other medicines taken with allopurinol?

Allopurinol may be taken in combination with other arthritis medicines, including:

- steroid medicines such as prednisolone or cortisone injections into the joint;
- anti-inflammatory medicines (NSAIDs) such as naproxen (Naprosyn) or ibuprofen (Brufen/Nurofen);
- colchicine (Colgout); and/or
- simple pain medicines such as paracetamol.

How long is the treatment continued?

Treatment is usually continued indefinitely as long as it is effective and as long as no serious side effects occur.

If allopurinol treatment is stopped suddenly, there is a high risk that your gout may worsen. It is very important not to stop your treatment unless advised by your doctor or unless side effects develop.

Are there any possible side effects?

Most people do not experience side effects from allopurinol. Below are possible side effects that you might experience with your treatment. Tell your doctor if you experience any side effects.

If you do experience side effects, a reduction in dose may minimise these so that you can continue to take the medicine. Your doctor will advise on any dose changes that are necessary.

Most common possible side effects:

- The most common side effects are *nausea or vomiting*. These can be reduced if you eat little and often. If you are sick, drink plenty of liquid.

Less common or rare possible side effects:

There are some rare but potentially serious side effects with allopurinol.

- ***Skin problems:*** Allopurinol can cause a rash or flaking skin, as well as boils or sore lips or mouth. **If any of these occur contact your doctor straight away.**
- ***Tiredness:*** Drowsiness can occur. If it makes you feel sleepy, avoid driving or operating machinery.
- ***Liver:*** Allopurinol can inflame the liver causing a type of hepatitis. Blood tests can pick this up if it occurs.
The dose of allopurinol may need to be reduced or it may need to be stopped if problems occur. Contact your doctor immediately if you notice yellowing of the skin and whites of the eyes.
- ***Other:*** Headache, dizziness, taste disturbances, high blood pressure, feeling generally unwell, and hair loss can occur.

Long term possible side effects:

Allopurinol can be taken for long periods to manage gout. There seem to be no long term side effects.

Allopurinol does not affect a person's ability to have children in the long term. See also ***Precautions.***

What precautions are necessary?

Blood tests:

- You may need to have blood tests during the first few months of treatment depending on what other medicines you are taking and on your other health concerns. The uric acid level in your blood will be checked to make sure the medicine is working.
- It is important to see your general practitioner regularly as they have an important role in monitoring your condition.

Other medicines:

- Allopurinol can interact with other medicines. You should tell your doctor (including your general practitioner, rheumatologist and others) about all medicines you are taking or plan to take. This includes over the counter or herbal/naturopathic medicines. You should also mention your treatment when you see other health professionals.
- **VERY SERIOUS INTERACTIONS will occur with azathioprine (Imuran) and mercaptopurine (Puri Nethol).**
- Other medicines that may interfere with allopurinol include: warfarin, ampicillin, amoxicillin, thiazide diuretics and sulfinpyrazone.
- Aspirin can be used safely in the low doses taken for prevention of heart attack and stroke. You should avoid taking aspirin in all other circumstance as at higher doses it can raise the uric acid level.

- Allopurinol can be taken safely with anti-inflammatory drugs (NSAIDs), as long as your kidney function is normal.
- The simple pain reliever, paracetamol, and combined medicines such as Panadeine and Panadeine Forte, can be used while taking allopurinol provided you take them as directed.

Alcohol:

- Alcohol can trigger an attack of gout. When taking allopurinol, keep your alcohol intake to a minimum i.e. 1-2 standard drinks, once or twice a week. Drinking more than 4 standard drinks on one occasion, even if infrequently, is strongly discouraged. In some cases total abstinence from alcohol is recommended. Check with your doctor about your situation.
- In addition to alcohol, other things that may trigger an acute gout attack include dehydration, diuretics (fluid tablets) and stopping allopurinol treatment.

Pregnancy and breastfeeding:

- The effects of allopurinol during pregnancy have not been well studied, so it is not clear if allopurinol causes birth defects.
- If you are pregnant or are considering having a child, you should discuss this with your doctor before beginning this medication.

All patients taking allopurinol should be seen regularly by a doctor to optimise treatment and to minimise any potential side effects. You must have regular blood tests when taking this medicine. Failure to do so will increase your risk of side effects. If you have any questions or concerns write them down and discuss them with your doctor.

Your doctor's contact details:

REMEMBER – Keep all medicines out of reach of children

The information in this sheet has been obtained from various sources and has been reviewed by the Australian Rheumatology Association. It is intended as an educational aid and does not cover all possible uses, actions, precautions, side effects, or interactions of the medicines mentioned. This information is not intended as medical advice for individual problems nor for making an individual assessment of the risks and benefits of taking a particular medicine. It can be reproduced in its entirety but cannot be altered without permission from the ARA.

The NHMRC publication: *How to present the evidence for consumers: preparation of consumer publications* (2000) was used as a guide in developing this publication.